

Atlantic Continental Shelf and Slope of the United States— Macrobenthic Invertebrate Fauna of the Middle Atlantic Bight Region—Faunal Composition and Quantitative Distribution

By ROLAND L. WIGLEY and ROGER B. THEROUX

GEOLOGICAL SURVEY PROFESSIONAL PAPER 529-N

*A description of the quantitative distribution
of macrobenthic invertebrate animals in
relation to geographic location, water
depth, bottom sediments, and range in
bottom water temperature*



UNITED STATES DEPARTMENT OF THE INTERIOR

JAMES G. WATT, *Secretary*

GEOLOGICAL SURVEY

Doyle G. Frederick, *Acting Director*

Library of Congress Cataloging in Publication Data (Revised)

United States. Geological Survey.

Atlantic Continental Shelf and slope of the United States.

(Geological Survey Professional paper; 529-N)

Program conducted jointly by the U. S. Geological Survey and the Woods Hole Oceanographic Institution.
Includes bibliographies.

CONTENTS: A. Geologic background, by K. O. Emery.—B. Physiography and sediments of the deep-sea basin, by R. M. Pratt.—[etc.]—N. Macrobenthic invertebrate fauna of the Middle Atlantic bight region, by R. L. Wigley and R. B. Theroux.

1. Continental margins—United States. 2. Oceanography—Atlantic coast (United States). 3. Marine biology—Atlantic coast (United States). I. Woods Hole, Mass. Oceanographic Institution. II. Title. III. Series.

QE75.P9 no. 529 557.3s [551.4'1] 79-604860

For sale by the Superintendent of Documents, U.S. Government Printing Office
Washington, D.C. 20402

CONTENTS

	Page		Page
Abstract	N1	Relation to bottom sediments	N124
Introduction	1	Distribution of sediment types	124
Reconnaissance survey	1	Total macrobenthic fauna of all taxonomic groups	124
Middle Atlantic Bight region	2	Entire Middle Atlantic Bight region	124
Previous studies	2	Subareas	132
Materials and methods	5	Taxonomic groups	137
Macrofauna samples	5	Entire Middle Atlantic Bight region	137
Benthos sampling gear	7	Subareas	137
Sample processing	7	Relation to sediment organic carbon	137
Data reduction	7	Distribution of sediment organic carbon	137
Bathymetry	10	Total macrobenthic fauna of all taxonomic groups	137
Temperature	10	Taxonomic groups	137
Geological samples	10	Entire Middle Atlantic Bight region	137
Faunal composition	10	Southern New England	154
Entire Middle Atlantic Bight region	10	Relation to range in bottom water temperature	154
Subarea differences in composition	17	Total macrobenthic fauna of all taxonomic groups	166
Geographic distribution	18	Entire Middle Atlantic Bight region	166
Total macrobenthic fauna of all taxonomic groups	18	Subareas	170
Major taxonomic components	23	Southern New England	170
Selected genera and species	95	New York Bight	170
Phylum Annelida	95	Chesapeake Bight	170
Phylum Pogonophoro	95	Taxonomic groups	170
Phylum Mollusca	95	Entire Middle Atlantic Bight region	170
Phylum Arthropoda	95	Subarea differences in distribution of	
Phylum Echinodermata	101	taxonomic groups	170
Bathymetric distribution	101	Dominant faunal components	192
Total macrobenthic fauna of all taxonomic groups	101	Bays and sounds	192
Entire Middle Atlantic Bight region	101	Continental Shelf	192
Subareas	106	Continental Slope	192
Taxonomic groups	108	Continental Rise	195
Entire Middle Atlantic Bight region	108	Acknowledgments	195
		References cited	195

ILLUSTRATIONS

		Page
FIGURE	1. Chart of the Middle Atlantic Bight region showing the location of geographical features and the three subarea divisions: Southern New England, New York Bight, and Chesapeake Bight ..	N3
	2. Chart showing station locations where quantitative samples of macrobenthic invertebrates were obtained	6
	3. Photograph showing side view of the Smith-McIntyre spring-loaded bottom sampler in the closed position	8
	4. Photograph showing bottom view of Campbell grab sampler	9
	5. Pie charts illustrating the taxonomic composition of the total macrobenthic fauna in the entire Middle Atlantic Bight region	12

	Page
FIGURE 6. Pie charts illustrating the taxonomic composition of the total macrobenthic fauna for each sub-area in the Middle Atlantic Bight region	N19
7-79. Maps of the Middle Atlantic Bight region showing the geographic distribution of:	
7. Density of all taxonomic groups combined	24
8. Biomass of all taxonomic groups combined	25
9. Density of Porifera	26
10. Biomass of Porifera	27
11. Density of Coelenterata	28
12. Biomass of Coelenterata	29
13. Density of Hydrozoa	30
14. Biomass of Hydrozoa	31
15. Density of Alcyonaria	32
16. Biomass of Alcyonaria	33
17. Density of Zoantharia	34
18. Biomass of Zoantharia	35
19. Density of Platyhelminthes	37
20. Biomass of Platyhelminthes	38
21. Density of Nemertea	39
22. Biomass of Nemertea	40
23. Density of Nematoda	41
24. Biomass of Nematoda	42
25. Density of Annelida	43
26. Biomass of Annelida	44
27. Density of Pogonophora	45
28. Biomass of Pogonophora	46
29. Density of Sipuncula	47
30. Biomass of Sipuncula	48
31. Density of Echiura and Priapulida	49
32. Biomass of Echiura and Priapulida	50
33. Density of Mollusca	52
34. Biomass of Mollusca	53
35. Density of Cephalopoda and Polyplacophora	54
36. Biomass of Cephalopoda and Polyplacophora	55
37. Density of Gastropoda	56
38. Biomass of Gastropoda	57
39. Density of Bivalvia	58
40. Biomass of Bivalvia	59
41. Density of Scaphopoda	60
42. Biomass of Scaphopoda	61
43. Density of Arthropoda	62
44. Biomass of Arthropoda	63
45. Density of Arachnida, Copepoda, Nebaliacea, Ostracoda, and Pycnogonida	65
46. Density of Cirripedia	66
47. Biomass of Cirripedia	67
48. Density of Cumacea	68
49. Biomass of Cumacea	69
50. Density of Tanaidacea	70
51. Biomass of Tanaidacea	71
52. Density of Isopoda	72
53. Biomass of Isopoda	73
54. Density of Amphipoda	74
55. Biomass of Amphipoda	75
56. Density of Mysidacea	76
57. Biomass of Mysidacea	77
58. Density of Decapoda	78
59. Biomass of Decapoda	79
60. Density of Bryozoa and Brachiopoda	81
61. Biomass of Bryozoa and Brachiopoda	82

	Page
FIGURES 7-79. Maps of the Middle Atlantic Bight region showing the geographic distribution of—Continued	
62. Density of Echinodermata	N83
63. Biomass of Echinodermata	84
64. Density of Holothuroidea	85
65. Biomass of Holothuroidea	86
66. Density of Echinoidea	87
67. Biomass of Echinoidea	88
68. Density of Ophiuroidea	89
69. Biomass of Ophiuroidea	90
70. Density of Asteroidea	91
71. Biomass of Asteroidea	92
72. Density of Ascidiacea and Hemichordata	93
73. Biomass of Ascidiacea and Hemichordata	94
74. Three selected species of Annelida and one Pogonophora	96
75. Selected bivalves, phylum Mollusca	97
76. Selected bivalves and gastropods, phylum Mollusca	98
77. Selected amphipods, phylum Arthropoda	99
78. A selected isopod and decapods, phylum Arthropoda	100
79. Selected echinoids, asteroids, and ophiuroids, phylum Echinodermata	102
80. Graph showing relationship between number of individuals and water depth for each subarea and for the entire Middle Atlantic Bight region	103
81. Graph showing relationship between biomass (wet weight) and water depth for each subarea and for the entire Middle Atlantic Bight region	104
82-87. Graphs showing density and biomass in relation to water depth in the entire Middle Atlantic Bight region for:	
82. Porifera, Hydrozoa, Alcyonaria, Zoantharia, Platyhelminthes, and Nemertea	110
83. Nematoda, Annelida, Pogonophora, Sipuncula, Echiura, and Priapulida	111
84. Polyplacophora, Gastropoda, Bivalvia, Scaphopoda, Cephalopoda, and Pycnogonida	112
85. Ostracoda, Cirripedia, Copepoda, Nebaliacea, Cumacea, and Tanaidacea	113
86. Isopoda, Amphipoda, Mysidacea, Decapoda, Bryozoa, and Brachiopoda	114
87. Holothuroidea, Echinoidea, Ophiuroidea, Asteroidea, Hemichordata, and Ascidiacea	115
88. Map showing geographic distribution of bottom-sediment types in the Middle Atlantic Bight region	125
89-94. Photographs showing:	
89. Gravel bottom at a depth of 23 m in the Nantucket Shoals region, south of Cape Cod, Mass	126
90. Sand bottom, containing small amounts of shell, on the Continental Shelf northeast of Cape Charles, Va., at a depth of 48 m	127
91. Silty-sand bottom at a depth of 406 m on the Continental Slope east of New Jersey	128
92. Sand bottom inhabited by a dense assemblage of sand dollars (<i>Echinarachnius parma</i>) at a depth of 48 m near midshelf east of Delaware	129
93. Sand-shell bottom at a depth of 69 m near the Outer Continental Shelf northeast of Cape May, N. J.	130
94. Silty-sand bottom at a depth of 178 m on the Outer Continental Shelf near Hudson Channel, south of New York City	131
95-104. Graphs showing:	
95. Relation between number of individuals and bottom-sediment types for the entire Middle Atlantic Bight region	133
96. Relation between biomass and bottom-sediment types for the entire Middle Atlantic Bight region	134
97. Relation between number of individuals and bottom-sediment types for each subarea	135
98. Relation between biomass and bottom-sediment types for each subarea	136
99. Density and biomass in relation to bottom sediments in the entire Middle Atlantic Bight region for Porifera, Hydrozoa, Alcyonaria, Zoantharia, Platyhelminthes, and Nemertea	139

	Page
FIGURES 95-104. Graphs showing—Continued	
100. Density and biomass in relation to bottom sediments in the entire Middle Atlantic Bight region for Nematoda, Annelida, Pogonophora, Sipuncula, Echiura, and Priapulida	N140
101. Density and biomass in relation to bottom sediments in the entire Middle Atlantic Bight region for Polyplacophora, Gastropoda, Bivalvia, Scaphopoda, Cephalopoda, Pycnogonida	141
102. Density and biomass in relation to bottom sediments in the entire Middle Atlantic Bight region for Ostracoda, Cirripedia, Copepoda, Nebaliacea, Cumacea, and Tanaidacea ..	143
103. Density and biomass in relation to bottom sediments in the entire Middle Atlantic Bight region for Isopoda, Amphipoda, Mysidacea, Decapoda, Bryozoa, and Brachiopoda ..	144
104. Density and biomass in relation to bottom sediments in the entire Middle Atlantic Bight region for Holothuroidea, Echinoidea, Ophiuroidea, Asteroidea, Hemichordata, and Ascidiacea	145
105. Map showing the geographic distribution of organic carbon in the bottom sediments of the Middle Atlantic Bight region	152
106. Graph showing the relation between number of individuals and sediment organic carbon for each subarea and for the entire Middle Atlantic Bight region	153
107. Graph showing the relation between biomass and sediment organic carbon for each subarea and for the entire Middle Atlantic Bight region	159
108-113. Graphs showing density and biomass in relation to sediment organic carbon in the entire Middle Atlantic Bight region for:	
108. Porifera, Hydrozoa, Alcyonaria, Zoantharia, Platyhelminthes, and Nemertea	160
109. Nematoda, Annelida, Pogonophora, Sipuncula, Echiura, and Priapulida	161
110. Polyplacophora, Gastropoda, Bivalvia, Scaphopoda, Cephalopoda, and Pycnogonida	162
111. Ostracoda, Cirripedia, Copepoda, Nebaliacea, Cumacea, and Tanaidacea	163
112. Isopoda, Amphipoda, Mysidacea, Decapoda, Bryozoa, and Brachiopoda	164
113. Holothuroidea, Echinoidea, Ophiuroidea, Asteroidea, Hemichordata, and Ascidiacea	165
114. Map showing distribution of the range in bottom-water temperature for the Middle Atlantic Bight region	167
115. Graph showing relation between number of individuals and range in bottom-water temperature for each subarea and for the entire Middle Atlantic Bight region	168
116. Graph showing relation between biomass and range in bottom-water temperature for each subarea and for the entire Middle Atlantic Bight region	169
117-122. Graphs showing density and biomass in relation to range in bottom-water temperature in the entire Middle Atlantic Bight region for:	
117. Porifera, Hydrozoa, Alcyonaria, Zoantharia, Platyhelminthes, and Nemertea	173
118. Nematoda, Annelida, Pogonophora, Sipuncula, Echiura, and Priapulida	174
119. Polyplacophora, Gastropoda, Bivalvia, Scaphopoda, Cephalopoda, and Pycnogonida	175
120. Ostracoda, Cirripedia, Copepoda, Nebaliacea, Cumacea, and Tanaidacea	176
121. Isopoda, Amphipoda, Mysidacea, Decapoda, Bryozoa, and Brachiopoda	177
122. Holothuroidea, Echinoidea, Ophiuroidea, Asteroidea, Hemichordata, and Ascidiacea	178
123. Map showing the geographic distribution of the number of individuals for each dominant taxon in the entire Middle Atlantic Bight region	193
124. Map showing the geographic distribution of the biomass for each dominant taxon in the entire Middle Atlantic Bight region	194

TABLES

	Page
TABLE 1. Research vessels, cruise identification and dates, and number of stations sampled	N5
2. Areas of several bathymetric zones within each subarea and total area of Middle Atlantic Bight region ..	5
3. Quantitative taxonomic composition of the macrobenthic invertebrate fauna, in both number of individuals and biomass, representing the entire Middle Atlantic Bight region	11
4. Invertebrate species contained in quantitative samples taken within the Middle Atlantic Bight region ..	13
5. Quantitative taxonomic composition of the macrobenthic invertebrate fauna, in both number of individuals and biomass, representing the Southern New England subarea	20
6. Quantitative taxonomic composition of the macrobenthic invertebrate fauna, in both number of individuals and biomass, representing the New York Bight subarea	21
7. Quantitative taxonomic composition of the macrobenthic invertebrate fauna, in both number of individuals and biomass, representing the Chesapeake Bight subarea	22
8. Number of samples within each depth range class in each subarea and for Middle Atlantic Bight region ..	101
9. Mean number of individuals and biomass of the macrobenthic invertebrate fauna in relation to water depth for each subarea and for the entire Middle Atlantic Bight region	101
10. Change and rate of change in density of invertebrates in relation to water depth	105
11. Change and rate of change in biomass of invertebrates in relation to water depth	105
12. Mean number of individuals listed by major taxonomic groups for each bathymetric class, representing the entire Middle Atlantic Bight region	108
13. Mean biomass listed by major taxonomic groups for each bathymetric class, representing the entire Middle Atlantic Bight region	109
14. Mean number of individuals listed by major taxonomic groups for each bathymetric class, representing the Southern New England subarea	116
15. Mean biomass listed by major taxonomic groups for each bathymetric class, representing the Southern New England subarea	117
16. Mean number of individuals listed by major taxonomic groups for each bathymetric class, representing the New York Bight subarea	118
17. Mean biomass listed by major taxonomic groups for each bathymetric class, representing the New York Bight subarea	119
18. Mean number of individuals listed by major taxonomic groups for each bathymetric class, representing the Chesapeake Bight subarea	121
19. Mean biomass listed by major taxonomic groups for each bathymetric class, representing the Chesapeake Bight subarea	122
20. Number of samples for each bottom sediment type in each subarea and for the entire Middle Atlantic Bight region	124
21. Mean number of individuals and biomass of the macrobenthic invertebrate fauna in relation to bottom sediments for each subarea and for the entire Middle Atlantic Bight region	132
22. Mean number of individuals listed by taxonomic groups in each bottom-sediment type for the entire Middle Atlantic Bight region	138
23. Mean biomass of each taxonomic group listed by bottom-sediment type for the entire Middle Atlantic Bight region	139
24. Mean number of individuals listed by taxonomic group in each bottom-sediment type for the Southern New England subarea	146
25. Mean biomass of each taxonomic group listed by bottom-sediment type for the Southern New England subarea	147
26. Mean number of individuals listed by taxonomic group in each bottom-sediment type for the New York Bight subarea	148
27. Mean biomass of each taxonomic group listed by bottom-sediment type for the New York Bight subarea ..	149
28. Mean number of individuals listed by taxonomic group in each bottom-sediment type for the Chesapeake Bight subarea	150
29. Mean biomass of each taxonomic group listed by bottom-sediment type in the Chesapeake Bight subarea ..	151
30. Number of samples for each class of sediment organic carbon in each subarea and for the entire Middle Atlantic Bight region	154

	Page
TABLE 31. Mean number of individuals and biomass of the macrobenthic invertebrate fauna in relation to percent organic carbon in bottom sediments for each subarea and for the entire Middle Atlantic Bight region --	N154
32. Mean number of individuals of each taxonomic group listed by sediment organic carbon content class, representing the entire Middle Atlantic Bight region -----	155
33. Mean biomass of each taxonomic group listed by sediment organic carbon content class, representing the entire Middle Atlantic Bight region -----	156
34. Mean number of individuals of each taxonomic group listed by sediment organic carbon content class, representing the Southern New England subarea -----	157
35. Mean biomass of each taxonomic group listed by sediment organic carbon content class, representing the Southern New England subarea -----	158
36. Number of samples within each water temperature range class in each subarea and for the Middle Atlantic Bight region -----	166
37. Mean number of individuals and biomass of the macrobenthic invertebrate fauna, all taxonomic groups combined, in relation to range in bottom-water temperature -----	166
38. Mean number of individuals of each taxonomic group listed by temperature-range class, representing the entire Middle Atlantic Bight region -----	171
39. Mean biomass of each taxonomic group listed by temperature-range class, representing the entire Middle Atlantic Bight region -----	172
40. Mean number of individuals of each taxonomic group listed by temperature-range class, representing the Southern New England subarea -----	180
41. Mean number of individuals of each taxonomic group listed by temperature-range class, representing the New York Bight subarea -----	181
42. Mean number of individuals of each taxonomic group listed by temperature-range class, representing the Chesapeake Bight subarea -----	182
43. Mean biomass of each taxonomic group listed by temperature-range class, representing the Southern New England subarea -----	183
44. Mean biomass of each taxonomic group listed by temperature-range class, representing the New York Bight subarea -----	184
45. Mean biomass of each taxonomic group listed by temperature-range class, representing the Chesapeake Bight subarea -----	185
